

Robotic-assisted early mobilization and virtual reality: a perspective on innovative support strategies for critically ill patients

Claudio Parco, Vanessa Kreuels, Malte Kelm, Christian Jung, Georg Wolff

Article - Version of Record



Suggested Citation:

Parco, C., Kreuels, V., Kelm, M., Jung, C., & Wolff, G. (2023). Robotic-assisted early mobilization and virtual reality: a perspective on innovative support strategies for critically ill patients. *Intensive Care Medicine Experimental*, 11, Article 86. <https://doi.org/10.1186/s40635-023-00571-x>

Wissen, wo das Wissen ist.

This version is available at:

URN: <https://nbn-resolving.org/urn:nbn:de:hbz:061-20241031-095036-5>

Terms of Use:

This work is licensed under the Creative Commons Attribution 4.0 International License.

For more information see: <https://creativecommons.org/licenses/by/4.0>

LETTERS TO THE EDITOR

Open Access



Robotic-assisted early mobilization and virtual reality: a perspective on innovative support strategies for critically ill patients

Claudio Parco¹, Vanessa Kreuels¹, Malte Kelm^{1,2}, Christian Jung^{1*} and Georg Wolff¹

Keywords Virtual reality, Robotic-assisted early mobilization, Intensive care

To the Editor

Survivors of critical illness frequently suffer from physical, psychosocial and cognitive impairments which result in a substantial loss in quality of life. Many research groups are working on structured rehabilitation programs to reduce the burden and there are preventive approaches like early mobilization, which innovative technical solutions may help to integrate into daily practice: virtual reality (VR) devices [1] and robotic-assisted early mobilization offer new and promising opportunities to improve patient rehabilitation on intensive care units (ICU).

VR can be adapted to various applications, including clinical scenarios ranging from patient education and staff training to support of conscious sedation in invasive procedures [2, 3]. VR interventions have shown efficacy and minimal side effects, which makes them a potential supportive therapeutic option. However, quality of evidence on the influence of VR-based interventions in critically ill patients is limited and randomized controlled trials are lacking.

Early mobilization is an integral part of modern critical care, which aims to improve functional capacity of patients after discharge. However, timing of conventional early mobilization is challenging for both patients as well as ICU staff [4], as it may be hindered by tubes and lines necessary for ventilation, feeding and intravenous medication of the patient in the early stages of recovery and may even provoke hazardous situations potentially resulting in adverse events [5]. Robotic systems were thus developed to support early mobilization (e.g., VEMOTION[®]) consisting of a specialized bed and the actual robot facilitate verticalization and a subsequent walking experience. Since the patient is fastened to the robot in the ICU bed without the need to transfer the patient to a separate machine, hazards and necessary manpower are potentially reduced with this method.

We have introduced a combination of both VR and robotic-assisted devices to facilitate early mobilization (Fig. 1): We initially applied this combination to support recovery of a 57-year-old patient, who was treated at our ICU for septic pneumonia. He entered a phase of prolonged ventilator weaning after initial stabilization. He enjoyed this combination and asked to use it several times a day for up to one hour. There were no adverse safety events during his treatment. This initial experience encouraged of to apply this combination to selected patients that potentially benefit from both the physical and psychological effects during their prolonged ICU treatment. We have included additional information about the patients in the Additional file 1. There is however room for further refinement since synchronization

*Correspondence:

Christian Jung
christian.jung@med.uni-duesseldorf.de

¹ Department of Cardiology, Pulmonology, and Vascular Medicine, University Hospital Düsseldorf, Medical Faculty of the Heinrich Heine University Düsseldorf, Moorenstr. 5, 40225 Düsseldorf, Germany

² Cardiovascular Research Institute Düsseldorf, Medical Faculty and University Hospital Düsseldorf, CARID, Heinrich-Heine-University, Düsseldorf, Germany



Fig. 1 A combination of Virtual Reality (e.g. Oculus Meta Quest 2[®]) and robotic-assisted early mobilization (e.g. VEMOTION[®] system, Reactive Robotics) in three critically ill patients (for more information about the patients see Additional file 1)

of the robotic system and VR headsets was technically impossible.

Conclusion

Prevention of physical and cognitive impairment following intensive care treatment requires innovative, multimodal therapeutic approaches: VR interventions and robotic-assisted early mobilization are easy to combine. However, there is a need for randomized clinical trials to further evaluate benefits and risks of these innovative methods while using standardized protocols and tools for outcome assessment. While application of this combination worked well in our selected patients, further research is necessary to prove feasibility in more challenging patients which e.g. suffer from delirium or motion sickness.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s40635-023-00571-x>.

Additional file 1. Background, patients and setting.

Author contributions

CP and GW wrote the manuscript and created media content. CJ initiated and supervised the project. VK and MK critically revised the manuscript. All authors consented to the manuscript for publication. The authors read and approved the final manuscript.

Funding

Virtual reality devices were funded by the State of North Rhine Westphalia (Giga for Health: 5GMedizin-campus. NRW, Project number 005-2008-0055 and PROFILNRW-2020-107-A, TP4)

Availability of data and materials

Data sharing not applicable to this article as no datasets were generated or analyzed during the current study.

Declarations

Ethical approval and consent to participate

Not applicable.

Consent for publication

The patient provided written consent to publication of his case and image material. Images are meant as examples for potential use. Local regulations for the different devices need to be carefully evaluated and instructions for use followed.

Competing interests

The authors declare that they have no competing interests.

Received: 21 August 2023 Accepted: 26 November 2023

Published online: 06 December 2023

References

1. Jung C et al (2022) Virtual and augmented reality in cardiovascular care: state-of-the-art and future perspectives. *JACC Cardiovasc Imaging* 15(3):519–532
2. Bruno RR et al (2020) Virtual reality-assisted conscious sedation during transcatheter aortic valve implantation: a randomised pilot study. *Eurointervention* 16(12):e1014–e1020
3. Bruno RR et al (2022) Virtual and augmented reality in critical care medicine: the patient's, clinician's, and researcher's perspective. *Crit Care* 26(1):326
4. Schweickert WD, Patel BK, Kress JP (2022) Timing of early mobilization to optimize outcomes in mechanically ventilated ICU patients. *Intensive Care Med* 48(10):1305–1307
5. TEAM Study Investigators and the ANZICS Clinical Trials Group (2022) Early active mobilization during mechanical ventilation in the ICU. *N Engl J Med*. <https://doi.org/10.1056/NEJMoa2209083>

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.