

Microcirculation information in clinical decision making: Rome wasn't built in a day

Raphael Romano Bruno, Glenn Hernandez, Jakob Wollborn, Bernd Saugel, Christian Jung on behalf of of the DAMIS study group

Article - Version of Record



Suggested Citation:

Bruno, R. R., Hernandez, G., Wollborn, J., Saugel, B., & Jung, C. (2023). Microcirculation information in clinical decision making: Rome wasn't built in a day. *Intensive Care Medicine*, 49(10), 1272–1273.
<https://doi.org/10.1007/s00134-023-07216-3>

Wissen, wo das Wissen ist.

This version is available at:

URN: <https://nbn-resolving.org/urn:nbn:de:hbz:061-20250226-103758-3>

Terms of Use:

This work is licensed under the Creative Commons Attribution 4.0 International License.

For more information see: <https://creativecommons.org/licenses/by/4.0>

CORRESPONDENCE



Microcirculation information in clinical decision making: Rome wasn't built in a day

Raphael Romano Bruno¹, Glenn Hernandez², Jakob Wollborn³, Bernd Saugel⁴ and Christian Jung^{1,5*}  on behalf of the DAMIS study group

© 2023 The Author(s)

We thank Edul and Dubin [1] and Damiani and colleagues [2] for their constructive criticism of our Direct Assessment of Microcirculation In Shock (DAMIS) trial [3]. DAMIS was a trial on the effect of integrating microcirculatory information into treatment considerations on mortality in patients with circulatory shock.

Direct visualization of the sublingual microcirculation using handheld video microscopy is used for research for more than 20 years. Recording and analyzing video sequences is still cumbersome, but time-consuming. To integrate sublingual microcirculation monitoring in clinical decision-making, it needs to be available at the bedside and provide microcirculatory variables in real time [4]. We, therefore, used the AVA 4.3C analysis software that allows automated, fast, and user-independent analysis of the sublingual microcirculation [5, 6]. Although the AVA 4.3C analysis software is not validated against manual gold standard analyses, we performed all measurements according to the manufacturer's instructions and current guideline recommendations [7].

We used the proportion of perfused small vessels (sPPV) as target variable because it is easy to interpret (also for clinicians not experienced with microcirculatory analyses) and has been shown to predict hospital mortality [8]. The sPPV risk categories were chosen based on the literature and the manufacturer's instructions.

The DAMIS trial indeed included patients with different types of circulatory shock. The finding that considering sublingual microcirculation monitoring during treatment decisions was consistent across patients with all types of shock. It is a limitation of the trial that we did not use specific treatment protocols for different types of shock, but there is insufficient evidence for specific algorithms including microcirculatory values. Additionally, using a strict treatment protocol based on the results of an experimental device was impossible.

Regarding the timing of microcirculatory assessments in DAMIS, we repeated the initial measurements after a 24-h interval. More frequent measurements might have been desirable, but our study design reflects a practical approach for daily clinical application.

There were no differences in microcirculatory variables—including capillary refill time—between survivors and non-survivors. One reason may be that all measurements were performed after the immediate initial resuscitation. Additionally, patients often died because life-sustaining care was withdrawn—and not from initial shock. In fact, limiting life-sustaining therapy was a significant predictor of mortality, albeit not remarkably higher than in other studies [9]. It is important to distinctively report withdrawal of life-sustaining therapy a cause of death.

We naturally agree that it is the treatment and not the monitoring that determines patient outcomes. Our trial suggests that considering microcirculatory variables using AVA 4.3C during clinical decision-making for treatment optimization does not improve outcomes in patients with shock. Future research needs to determine different strategies to employ bedside assessed microcirculation into clinical decision-making.

*Correspondence: christian.jung@med.uni-duesseldorf.de

¹ Medical Faculty, Department of Cardiology, Pulmonology and Vascular Medicine, Heinrich-Heine-University Duesseldorf, Moorenstraße 5, 40225 Duesseldorf, Germany

Full author information is available at the end of the article

This comment refers to the article available online at <https://doi.org/10.1007/s00134-023-07098-5>.

Author details

¹ Medical Faculty, Department of Cardiology, Pulmonology and Vascular Medicine, Heinrich-Heine-University Duesseldorf, Moorenstraße 5, 40225 Duesseldorf, Germany. ² Departamento de Medicina Intensiva, Facultad de Medicina, Pontificia Universidad Católica de Chile, Santiago, Chile. ³ Department of Anesthesiology, Perioperative and Pain Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, USA. ⁴ Department of Anesthesiology, Center of Anesthesiology and Intensive Care Medicine, University Medical Center Hamburg-Eppendorf, Hamburg, Germany. ⁵ CARID (Cardiovascular Research Institute Düsseldorf), Duesseldorf, Germany.

Funding

Open Access funding enabled and organized by Projekt DEAL. This work was supported by the Forschungskommission of the Medical Faculty of the Heinrich-Heine-University Düsseldorf No. 2020-21 to RRB for a Clinician Scientist Track. Furthermore, institutional support has been received by the German Research Council (SFB 1116, B06) as well as the State of North Rhine Westphalia (Giga for Health: 5GMedizincampus. NRW, Project number 005-2008-0055 and PROFILNRW-2020-107-A, TP4). No (industry) sponsorship has been received for this investigator-initiated study.

Availability of data and material

The anonymized data can be requested from the authors if required.

Declarations

Conflicts of interest

The authors declare that they have no competing interests.

Ethics approval and consent to participate

The primary competent ethics committee was the Ethics Committee of the University of Duesseldorf, Germany. Institutional research ethic board approval was obtained from each study site.

Consent for publication

The manuscript does not contain any individual person's data in any form.

Open Access

This article is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License, which permits any non-commercial use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The

images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc/4.0/>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Accepted: 27 August 2023

Published: 11 September 2023

References

1. Kanoore Edul VS, Dubin A (2023) Pitfalls in the use of microcirculation as a resuscitation goal. *Intensive Care Med* <https://doi.org/10.1007/s00134-023-07191-9>
2. Damiani E, Scorzella C, Carsetti A, Donati A, Adrario E (2023) Microcirculation as a guide for therapy: do not condemn an innocent without a fair trial. *Intensive Care Med*. <https://doi.org/10.1007/s00134-023-07192-8>
3. Bruno RR et al (2023) Direct assessment of microcirculation in shock: a randomized-controlled multicenter study. *Intensive Care Med* 49(6):645–655
4. Monnet X, Saugel B (2018) Could resuscitation be based on microcirculation data? We are not sure. *Intensive Care Med* 44(6):950–953
5. Bruno RR et al (2020) Evaluation of a shorter algorithm in an automated analysis of sublingual microcirculation. *Clin Hemorheol Microcirc* 76(2):287–297
6. Bruno RR et al (2020) Sublingual microcirculation in prehospital critical care medicine: a proof-of-concept study. *Microcirculation* 27(5):e12614
7. Ince C et al (2018) Second consensus on the assessment of sublingual microcirculation in critically ill patients: results from a task force of the European Society of Intensive Care Medicine. *Intensive Care Med* 44(3):281–299
8. Spanos A et al (2010) Early microvascular changes in sepsis and severe sepsis. *Shock* 33(4):387–391
9. Bruno RR et al (2021) Management and outcomes in critically ill nonagenarian versus octogenarian patients. *BMC Geriatr* 21(1):576